

Ticks Found on Dogs

- Treat every dog in household or on property with long-acting tick prevention. Recommended products include:
 - Bravecto chewable (3 months)
 - Nexgard chewable (1 month)
 - Simparica or Simparica Trio chewable (1 month)
 - Seresto Collar (8 months)
 - K9 Advantix II (topical, 1 month)
 - Frontline Plus or Frontline Gold (topical, 1 month)
- Continue treatment year-round. Do not allow to lapse.

Ticks Found in the Yard or Home

- Treat any dogs in the home or on the property (see above)
- Use labelled pesticide on areas where ticks are present
 - Bifen IT (bifenthrin)
 - Bayer Advanced Multi-insect Killer (Beta-cyfluthrin and imidacloprid)
- Seal cracks in cement and other surfaces
- Yard cleanup
 - Remove outdoor furniture
 - Clean up woodpiles and other materials that ticks can hide under
- Recheck in 3 weeks
 - Place dry ice traps in areas where ticks were seen for 2-3 hours
 - Or examine dogs present
 - If ticks are present, repeat treatment
- Recheck again in 3 weeks

Notes on Environmental Treatment:

ALWAYS follow label instructions for application and personal protective equipment use when applying pesticides

- Liquid may be more effective than granules but increase risk of exposure to the applicator
- Typically only last 3 weeks at the most
- Brown dog ticks may be resistant to permethrin-based products (e.g., deltamethrin)

Tick or Ticks Found on Person

- Identify tick. If a brown dog tick, proceed with following steps.
 - If tick is not a brown dog tick, indoor treatment is not required
 - If tick is a brown dog tick, indoor and outdoor treatment may be needed
- Check dogs and other pets for ticks.
 - Treat dogs as described above.
- Check home and yard for ticks. If found, treat as above.
 - Check under yard debris, cracks of furniture, etc.